

Brief intervention on the consumption of fruits and vegetables and physical activity in university students

Clara Helena GONZÁLEZ CORREA¹, Aida María GONZÁLEZ CORREA¹, Diana María MUÑOZ PÉREZ²,
Carmen DUSSAN LUBERTH³

1 Faculty of Health Sciences, Basic Science Department, Universidad de Caldas.

2 Faculty of Health Sciences, Basic Science Department, Universidad Libre de Colombia.

3 Faculty of Exact Sciences, Department of Mathematics, Universidad de Caldas.

Recibido: 20/septiembre/2021. Aceptado: 28/noviembrebre/2021.

ABSTRACT

Introduction: The World Health Organization (WHO) has estimated that up to 2.7 million lives could be saved annually if enough fruits and vegetables were consumed. Regular physical activity has been shown to reduce the risk of various diseases and help with energy balance and weight control. For university students it is complex to follow healthy guidelines such as the consumption of fruits and vegetables and the realization of physical activity due to the lack of time to cook and academic commitments.

Objective: To determine the nutritional status and physical activity of university students before and after an educational intervention with WHO recommendations.

Materials and methods: A prospective longitudinal study in which 70 university students were evaluated. The intervention consisted of an individualized educational talk. The nutritional status was evaluated by means of a survey on the consumption of fruits and vegetables and anthropometric measurements and physical condition by recording the time of habitual physical activity per week and the Harvard step test. The data were analyzed using the statistical software XLSTAT version 2014.3.07.

Results and discussion: The intervention had a positive effect on healthy lifestyle habits, increasing the percentage of compliance with the physical activity recommendations, 16%

in men and 9% in women. In addition, the intake of fruits and vegetables increased by 14% and 12%, respectively.

Conclusions: These improvements provide a basis for designing and implementing strategies that increase the percentage of adherence to healthy lifestyle habits through low-cost and easy-to-adopt interventions.

KEY WORDS

Eating habits, Fruit and Vegetables, Physical Activity, University Students.

ABBREVIATURES

WHO: World Health Organization.

NCDs: Non- Communicable Chronic Diseases.

BMI: Body Mass Index.

WHR: Waist/Hip Ratio.

HGST: Hand Grip Strength Test.

BI: Brohwa Index.

PCA: Principal Component Analysis.

FVC: Fruits and Vegetables Consumption.

EXC: Exercise Time.

BC: Body Composition.

SD: Standard Deviation.

INTRODUCTION

It is estimated that 77% of deaths from non- communicable chronic diseases (NCDs) occur in low- and middle-income countries. Chronic diseases include heart disease, strokes,

Correspondencia:
Clara Helena GONZÁLEZ CORREA
clara.gonzalez@ucaldas.edu.co

cancer, chronic respiratory diseases, and diabetes mellitus. These diseases are, in large part, the result of an inadequate diet, physical inactivity, tobacco consumption and excess of alcohol¹. Low consumption of plant-based foods, including fruits and vegetables, is associated with an increased risk of several human NCDs. A 2018 study suggested that 12% lung disease, 19% gastric cancer, 19% stroke and 31% of ischemic heart disease globally could be avoided by increasing the daily intake of fruits and vegetables to at least 400 g per day². These facts generate a high economic burden, as well as social dependence and physical disability. This generates a high economic burden, as well as social dependence and physical disability in people of these countries³.

It has been shown that regular physical activity reduces the risk of coronary heart disease and stroke, type II diabetes, hypertension, colon cancer, breast cancer and depression, and helps to achieve energy balance and weight control⁴. On the other hand, adequate consumption of fruit and vegetables can prevent micronutrient deficiency and reduce the risk of heart disease, obesity, diabetes and cancer, specifically gastrointestinal cancer. To encourage the incorporation of physical activity and fruit and vegetable consumption as part of a healthy lifestyle, the WHO adopted the global strategy on diet, physical activity and health for the prevention and control of NCDs⁵. The goal for physical activity in adults aged between 18 and 64 was estimated minimum of 150 minutes of moderate aerobic physical activity, or 75 minutes of vigorous aerobic physical activity each week or an equivalent combination of moderate and vigorous activities⁴. As for the consumption of fruit and vegetables, the goal for the population over two years old is an intake of at least 400g of these foods each day, which equals five servings, for example three of fruit and two of vegetables⁵.

Adopting healthy habits is essential to maintain good health and avoid the risk of those diseases⁶. For university students, many of whom come from different cities in the country with diverse eating habits, it is complex to follow healthy guidelines. According to them, the lack of time to cook, eat or prepare healthy foods. In addition, the change from a regular routine of physical activity at school level to a routine of physical inactivity and full-time academic commitments, make it difficult to comply with the recommendations⁷.

OBJECTIVE

The objective of this study was to determine the nutritional status and physical activity of students benefiting from the food subsidy from a University in Manizales and carry out an educational intervention in compliance with the WHO recommendations related to physical activity and consumption of fruit and vegetables.

MATERIAL AND METHODS

From 2014 to 2015 a prospective longitudinal study was conducted, with volunteers from different faculties of a

University in Manizales, Colombia. In the first assessment, 103 students were examined. The criteria to be included in the study were being an active student in this institution, being between 18 and 40 years old and receiving nutritional support with a lunch from Monday to Friday. The exclusion criteria were being pregnant or having any physical condition that prevented the potential participants from undergoing the tests. The procedures were approved by the ethics committee of the Faculty of Health. The intervention was carried out with 30 female and 40 male students who completed the second assessments.

The measurements were taken in the nutritional assessment laboratory under previously standardized environmental procedures and conditions. Anthropometry was carried out according to the protocol from the American Society of Hand Therapists^{8,9} and physical condition using the Harvard step test. Waist and hip circumferences, weight, height, hand grip strength and physical condition were measured. From these data, Body Mass Index (BMI) (kg/m²), waist/hip ratio (WHR), Hand Grip Strength Test (HGST) (kg/f) and Brohua Index (BI) were calculated¹⁰. Total body fat percentage (% body fat) was obtained from four folds: bicipital, tricipital, subscapular and suprailiac, applying the formula of Durnin and Womersley, with Skyndex II® digital caliper that has the incorporated formula. After taking the measurements, to know the intake of the respondents, the questionnaire on the frequency of consumption was applied. Individuals reported their daily or weekly consumption of fruits and vegetables in terms of servings after the meaning of a serving of fruit or vegetables was explained to them. Each person received their results in a pre-coded format, and they were interpreted according to the WHO reference values for the BMI¹¹, % body fat¹², WHR by Colombian government¹³, HGST¹⁴ and BI according to the correspondent equation (BI= Duration of exercise in seconds x 100/2 x sum of pulse counts in recovery) and the following table (Table 1) for interpretation¹⁵.

Table 1. Interpretation of the Brohua Index results

Rating	Fitness Index
Excellent	> 90
Good	80-89
High Average	65-79
High Average	65-79
Low Average	55-64
Poor	< 55

Intervention

The intervention was an individualized face to face individual educational talk for 30 minutes. It consisted of explaining each person's results to emphasize their own deficits and needs. Their results were interpreted according to reference values described in methods section.

Everyone was instructed on the basic recommendations of a healthy lifestyle according to WHO⁵ and five very detailed reasons why it is important to consume at least five daily servings of fruits and vegetables: water soluble vitamins, minerals, water, fiber, and polyphenols content. They were also instructed on the need to diversify their consumption considering the meaning of the different colors and the antioxidant, anti-inflammatory and anti-neoplastic power that they represent. In addition, they were informed about the positive effects related to a higher intake of fruits and vegetables. Because so many students argued that fruits were expensive, a comparison was made between the price of a fruit such as banana or guava and a package of potato chips which can be up to five to six times more expensive in this low-to-middle income country.

Based on the physical activity guidelines for Americans, 2nd edition, the participants were also instructed to get a minimum of 150 minutes per week of moderate aerobic activity or 75 minutes per week of vigorous aerobic activity for two days a week¹⁵. Additionally, each student was explained the need to combine the above with a resistance muscle strengthening activity with weights twice a week. Emphasis was placed on the scientific evidence that shows a very close relationship between physical activity and immunological, physical, and mental well-being. They were also asked how they were transported to the university and encouraged to go to the university on foot. They were also encouraged to go down a few blocks before arriving at their destination so that if they used public transportation, they would walk at least a couple of blocks, and use every opportunity to walk.

Those who had their measurements in normal ranges and complied with the recommendations were congratulated and invited to continue their good habits. People with one or more of the examined parameters altered, were explained the long-term consequences especially related to NCDs, academic performance and cognitive impairment.

Participants Follow-Up

After one year, the same above measurements were made and a survey was performed to establish the effect of the intervention performed the previous year. The questionnaire included questions about consumption of fruit and vegetables, amount of physical activity performed before and after the intervention and perceived changes when they had a variation in their habits.

Statistical Analysis

The data were analyzed using the statistical software XLSTAT version 2014.3.07. Data were expressed as means and standard deviations. For the comparison of means, before and after the educational intervention, the student's t-test was applied for paired samples. For computing the statistical significance of the hypotheses, a two-tailed test was developed. A significance of 5% was used. In addition, the data were analyzed by principal component analysis (PCA) for men and women independently.

RESULTS

Seventy people finally participated in the study (30 female/40 male), with an average age of 21 (3) and 22 (4) and a BMI of 22.05 (3.5) and 21.9 (2.6) kg/m for women and men, respectively. During the second evaluation of fitness test, four women and one man had constraints which prevented them from being involved. The characteristics of the participants at the beginning and at the end of the study are reported in Table 2.

Significant favorable changes were observed in all variables except the BI for both sexes, % body fat in women and WHR in men. The results obtained show that, on average, both men and women had a normal BMI. However, women showed high values (>30%) for % body fat, while men had acceptable values (≤20%). Both, men, and women did not reach the percentage of strength (>85%) referenced for their age and sex and the WHR showed that women were at the upper limit of reference, while men were below it.

Regarding physical condition, the BI showed that women had a physical status classified as "good" while men were classified in the "average" category. Exercise time increased after the intervention in both women and men; however, only 26% of men complied with the recommendations of physical activity of the WHO. The average for all men was of 68 minutes per week (45.3% of the recommended level). After the intervention, 42% of participants already fulfilled the recommendation, raising the average physical activity to 98 minutes per week (65.3% of the recommended level). Likewise, before the intervention, only 20% of women complied with the activity recommendations and the average for this group was of 54 minutes per week (36% of the recommended level). After the intervention, 29% of them already fulfilled the recommendation, raising the average physical activity to 77 minutes per week (51.3% of the recommended level) (Figure. 1).

Regarding the consumption of fruit and vegetables, the average consumption at the beginning of the study for all participants was 1.3 portions and at the end, 2.0 and 1.8 for men and women, respectively. This reflects an increase in consumption of 53% and 38.4% respectively. Despite this, after the intervention, only 12% of women and 14% of men consumed the five servings recommended by WHO (Figure 1).

Students who followed at least one of the WHO recommendations noted some physical and emotional changes. In women


Table 2. Anthropometric Parameters, Fruit and Vegetables Consumption and Exercise Time Before and After the Intervention

	Women n=30			Men n=40		
	Before	After	p-value	Before	After	p-value
FVC (# of portions)	1.3 (1.1)	2.0 (0.9)	0.01	1.3 (0.9)	1.8 (1.0)	0.01
EXC (min)	40.4 (50.6)	146.0 (65.4)	0.00	75.8 (71.0)	162.3 (58.8)	0.00
Weight (kg)	54.4 (8.8)	57.4 (8.5)	0.00	63.2 (8.2)	68.5 (8.6)	0.00
BMI (Kg/m ²)	22.1(3.5)	23.1(3.2)	0.00	22.0 (2.7)	23.7 (2.8)	0.00
Body fat (%)	31.3 (3.6)	30.6(2.1)	0.21	18.9 (4.4)	20.0 (3.7)	0.01
WHR	0.7 (0.1)	0.8 (0.1)	0.04	0.8 (0.1)	0.8 (0.1)	0.50
HGST (Kg/f)	23.9 (3.9)	25.8 (4.6)	0.00	35.2 (6.0)	41.7 (6.3)	0.00
BI	78.0 (21.0)	73.2 (32.7)	0.38	73.9 (30.7)	71.0 (23.6)	0.24

Abbreviations: Fruits and vegetables consumption (FVC); Exercise Time (EXC); Body Mass Index (BMI); Waist/Hip Ratio (WHR); Hand Grip Step Test (HGST); Brohua Index (BI); Body Composition (BC).

For the comparison of means, before and after the educational intervention, the student's t-test was applied for paired samples. A significance of 5% was used.

Figure 1. Changes in exercise time per week and in the consumption of fruit in men and women



there was improvement in the perception of physical and mood well-being. 28% of female respondents said they felt an increase in energy, 16% an improvement in their physical condition, 15% noticed some loss of weight, 14% improved digestion, 7% the appearance of the skin, 4% increased concentration and 3% in the quality of sleep. Additionally, 5% reported decrease in stress. The men reported improvement in the perception of physical and mood well-being. 30% of male respondents said they felt an increase in energy and 22% an improvement in their physical condition, 8% in their digestion, 9% in their strength, 4% in their concentration, 3% in sleep quality, 10% noticed an increase in muscle mass, 5% in weight increase and 4% reported a decrease in stress.

The data were subjected to a PCA by sex. Bartlett's sphericity test showed that there were differences between the variables ($p < 0.001$ for women and men), which allowed the PCA to be performed. Three main components had eigen values greater than 1.0, which were used for subsequent analysis. These components explained 66.9% and 64.4% of the variance for women and men respectively (Table 3).

For women, the first component explains the greater variance (33.23%), where BMI (34.78%), % body fat (23.6%) and exercise (15%) contributed positively to the component. In the second component, those who contributed the most in a positive way were fruit consumption (24.8%) and BI (36%), HWR contribution was (22.3%) in a negative way. For women, the first component was related to exercise time, the second with fruit consumption and the third with both. In the third component, fruit consumption and exercise time (49.6% and 20.4% respectively) contributed positively. For men, the first component contributed to 26.9% of the total variance explained, with the variables that most contributed to the component in a positive way being the BMI (40.4%), the fat % (19.8%) and the HWR (19.5%). In the second component were BI (33.8%), HGST (22.9%) and EXC (22.4%). In the third component, those that contributed the most were WHR (38.7%) in a positive way and HGST (31.5%) in a negative way. The first component was related to body composition, the second to exercise and fruit consumption and the third to HGST (Table 4).

Table 3. Total variance explained by the components

	Women			Men		
	EXC	FVC	FV and EXC	BC	FVC and EXC	HGST
Eigen value	2.326	1.251	1.107	1.883	1.608	1.019
Variability (%)	33.233	17.874	15.809	26.907	22.971	14.551
% accumulated	33.233	51.107	66.916	26.907	49.878	64.430

Abbreviations: Fruits and vegetables consumption (FVC); Exercise Time (EXC); Body Composition (BC); Hand Grip Step Test (HGST). Data are mean and Standard Deviation (SD).

Table 4. Factor loadings for the three components obtained

	Women			Men		
	EXC	FVC	FVCEXC	BC	FVCEXC	HGST
FVC (# of portions)	-0.004	0.557	0.741	-0.080	0.459	-0.212
EXC (min)	0.599	0.092	0.476	-0.014	0.600	0.142
BMI (Kg/m ²)	0.900	-0.161	-0.235	0.872	0.208	-0.212
Body fat (%)	0.742	0.340	-0.177	0.856	-0.214	-0.340
WHR	0.552	-0.529	0.307	0.605	0.193	0.628
HGST (Kg/f)	0.550	0.245	-0.200	-0.134	0.606	-0.566
BI	-0.004	0.672	-0.331	0.015	0.738	0.278

Abbreviations: Fruits and vegetables consumption (FVC); Exercise Time (EXC); Body Mass Index (BMI); Waist/Hip Ratio (WHR); Hand Grip Step Test (HGST); Brohwa Index (BI); Body Composition (BC).

Variables with a factor load > 0.4 were considered a good contribution to the component.


In the factorial plans constituted by components one and two for women and men, it was observed that in women there was a direct relationship between the consumption of fruit and vegetables and the BI after the intervention (the higher the consumption of fruit the more the BI improves), independent of body composition (BMI, HWR, % body fat) and HGST (Fig. 2).

For men, fruit and vegetables consumption and exercise showed a direct relationship with the BI and the HGST (higher consumption of fruit and exercise time improves HGST and BI), independent of body composition (BMI, HWR, % body fat) (Fig. 2).

According to the recommendations of the health authorities at the world.

The results showed that indeed, the majority indeed most of the students (74.5%) did not comply with the WHO recommendation since they only consumed between one and two servings of fruits and vegetables per day. This is consistent with studies^{13,22} in Chilean university students where they found that only 5% of their students reached the goal of five daily servings, while in the present study 2% did. On the contrary, a study conducted at Universidad del Valle and Universidad ICESI in Colombia, found that 15% of the students surveyed followed the recommendations²³.

Figure 2. Factorial drawings of axes 1 and 2. A Women, B Men. F, fruit; V, vegetables



DISCUSSION

Various strategies have been used to promote healthy eating and physical activity habits in university students such as taking advantage of public university restaurants, teaching courses to improve culinary skills, using a food pantry, or even offering a monetary incentive¹⁶⁻¹⁹. The effects are still unsatisfactory, and more alternatives are required to improve them^{20,21}.

This study wanted to take advantage of those obtained by the university students during an evaluation of the nutritional status of students who received a food subsidy to try to sensitize them to a greater consumption of fruits and vegetables and the performance of regular physical activity ac-

In addition to contributing to the prevention of NCDs, the consumption of fruit and vegetables exerts beneficial effects by providing micronutrients such as vitamins A, C, folic acid, and minerals, as well as the presence of dietary fiber and some photochemical that contribute to the maintenance of good health². The low intake of these foods has been associated with a decrease in manual prehensile force²⁴. This could be the reason why 90% of men and 84% of women did not reach the level of strength required at the start of the study.

In general, Colombian population has a low consumption of fruits and vegetables and a high consumption of refined cereals, fats, and sweets with the risk of gaining weight and generating vitamin and mineral deficiencies. Thus, in this

country, the highest prevalence of low vitamin A intake occurs in the university community, reaching 80%. Therefore, it is necessary to break down barriers that students mention and prevent them from fully complying with the proposed recommendations. Socioeconomic factors, influence of the eating behaviors from family and friends, lack of information regarding following a healthy diet and lack of time are the main reasons they argue¹⁹.

Despite the poor compliance with the recommendations for fruit and vegetables consumption by university students, there was a positive effect of the intervention with a change in the percentage from 2% to 13% of students who reached the goal in this aspect. However, this figure is much lower than in United States where 40% of the population manages to meet the proposed goals²⁵.

Twenty-six % of men and 20% of women said they were doing physical activity for at least 30 minutes, five times or more per week and this value is like that found by Olivares and Lera¹⁴ who report 25% in men and 16% in women. The data also resemble those of a study conducted in Colombia at Universities in Cali, Bogotá, Manizales and Tuluá, in which the authors concluded that only 22.2% of students perform physical activity. It was evidenced that men carried out a higher percentage of physical activity than women²⁰. This pattern is repeated in Chile where men performed 15% more physical activity than women. This behavior could be caused by the fact that, culturally, women have been assigned to care of family members, reducing the chances of having enough time for themselves²⁶.

Students who complied with the recommendations of physical activity perceived a physical and mental well-being, possibly associated with the release of neurotransmitters such as serotonin, dopamine and norepinephrine that have been associated with memory recovery and a better mood²⁷.

The results obtained in this study suggest that an educational intervention constitutes an effective action to achieve a better adherence of university students to the recommendations of a healthy lifestyle. It is probable that the individualized intervention, together with the interpretation of its results immediately after obtaining them, evidencing in many cases that the students had deficits their muscular strength, excess fat, or poor physical condition, was moderately effective in sensitizing the students towards healthier habits.

A study carried out with individuals aged between 18 and 24 through a course to promote fruit and vegetables consumption in 2006 produced a significant increase in the consumption of fruit and vegetables ($p < 0.005^{**}$) and a reduction ($p < 0.05^{**}$) of the consumption of chips²⁸. On the other hand, another study with university students determined that after two years, an educational intervention was not effective in promoting long-term physical in men, although in women there was an improvement in behavior related to physical ac-

tivity. The intervention consisted of offering a general health course to the students as well as a guidance through emails and phone calls²⁹. A systematic review published in 2002, with the aim of evaluating the effectiveness of a variety of approaches to increase physical activity (informative, behavioral, social, environmental, and political) concluded that there is sufficient evidence that physical education in school is effective in increasing physical activity levels and improving physical condition³⁰.

The study has some limitations. There was no control group to compare the results. This was since the students belonged to a university subgroup that received support from the University for belonging to low socioeconomic strata and all of them underwent the same intervention. This is also why the results can only be applied to this subgroup of people. However, these circumstances are similar in the country's state universities and the results could be a reference for taking similar actions to those in this study. At present, we continue to make these assessments with students entering university and the behavior in relation to diet and exercise has not changed and that is why we consider that the results are still valid and applicable even if it is a study developed in 2015.

CONCLUSION

Although the improvements in fruits and vegetables consumption and physical activity carried out in this study were discrete and are still far from complying with the recommendations proposed by WHO the results provide a basis for designing and carrying out strategies that increase the percentage of adherence to healthy lifestyle habits through low-cost interventions and easy application.

ACKNOWLEDGMENTS

This research was made possible thanks to the support of volunteers, some medical students, and the University Welfare Unit.

BIBLIOGRAFÍA

1. Organización Mundial de la Salud (OMS). Enfermedades no transmisibles [Internet]. OMS; 2021 [Consultado 2021 Nov 23] Disponible en: <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases>.
2. Duthie GG, et al. Effect of increasing fruit and vegetable intake by dietary intervention on nutritional biomarkers and attitudes to dietary change: a randomised trial. *Eur J Nutr*. 2018; 57(5):1855-1872.
3. Villamizar GC, Arias G, Salgar W, Cala M. Boletín del Observatorio en Salud. *Rev Salud Publica*. 2010;33(1):67-78.
4. Organización Mundial de la Salud (OMS). Recomendaciones Mundiales sobre la Actividad Física para la Salud. 2012;66:37-9.

5. Organización Mundial de la Salud (OMS). Actividad Física [Internet]. OMS; 2018 [Consultado 2021 Nov 23]. Disponible en: <https://www.who.int/news-room/facts-in-pictures/detail/physical-activity>
6. Troncoso PC, Amaya P. Factores sociales en las conductas alimentarias de estudiantes universitarios. *Rev Chil Nutr.* 2009;36(4):1090–7.
7. Espinoza LO, Rodríguez FR, Gálvez JC, Macmillan NK. Hábitos de alimentación y actividad física en estudiantes universitarios. *Rev Chil Nutr.* 2011;38(4):458–65.
8. Janssen IAN, Heymsfield SB, Baumgartner RN, Ross R, Heymsfield SB, Richard N, et al. Estimation of skeletal muscle mass by bioelectrical impedance analysis. 2000;465–71.
9. Roberts HC, Denison HJ, Martin HJ, Patel HP, Syddall H, Cooper C, et al. A review of the measurement of grip strength in clinical and epidemiological studies: Towards a standardised approach. *Age Ageing.* 2011;40(4):423–9.
10. Brohwa L, Health CW. Step tests method measuring physical fitness for hard muscular work in adult men. *Rev men. Rev Canadian Biol.* 1943; 2-86.
11. Benton D, Braun H, Cobo JC, Edmonds C, Elmadfa I, El-Sharkawy A, et al. Executive summary and conclusions from the European Hydration Institute expert conference on human hydration, health, and performance. *Nutr Rev.* 2015;73:148–50.
12. Gallagher D, Heymsfield SB, Heo M, Jebb SA, Murgatroyd PR, Sakamoto Y. Healthy percentage body fat ranges: An approach for developing guidelines based on body mass index. *Am J Clin Nutr.* 2000;72(3):694–701.
13. Olivares CS, Lera ML, Bustos ZN. Etapas del cambio, beneficios y barreras en actividad física y consumo de frutas y verduras en estudiantes universitarios de Santiago de Chile. *Rev Chil Nutr.* 2008;35(1):25–35.
14. Mathiowetz V, Kashman N, Volland G, Weber K, Dowe M, Rogers S. Grip, and pinch strength: Normative data for adults. *Arch. Phys. Med. Rehabil.* 1985; 66(2):69-74.
15. Hootman JM. "2008 Physical activity guidelines for americans: an opportunity for athletic 387 trainers". *Journal of Athletic Training.* 2009; 44(1): 5–6.
16. Hartmann Y, Botelho R, Akutsu R, Puppini ZR. Consumption of Fruits and Vegetables by Low-Income Brazilian Undergraduate Students: A Cross-Sectional Study *Nutrients.* 2018; 10(8):1121.
17. Nagatomo W, Saito J, Kondo N. Effectiveness of a low-value financial-incentive program for increasing vegetable-rich restaurant meal selection and reducing socioeconomic inequality: a cluster crossover trial. *The international journal of behavioral nutrition and physical activity.* 2019; 16(81): 1-10.
18. Greyce LB, Jomori MK, Fernandes AC, Colussi CF, Condrasky MD, et al. Positive impact of a cooking skills intervention among Brazilian university students: Six months follow-up of a randomized controlled trial. *Appetite.* 2018; 30: 247-255.
19. 20. Clerkin, K.D., Pohl, C.J., & Kim, M.J. Influencing nutritional habits of college students using a food pantry. *Journal of American College Health.* 2020; doi: 10.1080/07448481.2020.1721506 401.
20. 21. Siegel KR. Insufficient Consumption of Fruits and Vegetables among Individuals 15 Years and Older in 28 Low- and Middle-Income Countries: What Can Be Done? *J Nutr.* 2019; doi: 40310.1093/jn/nxz123.
21. 22. Darfour SA, Andrade JE, Grigsby DS. Review of policies to increase fruit and vegetable consumption and physical activity in 49 low- and middle-income countries. *Journal of Public Health.* 2019; 41 (1): 119– 407.
22. López-Espinoza MA. Etapas del cambio conductual ante la ingesta de frutas y verduras, control de peso y ejercicio físico de estudiantes de la Universidad del Desarrollo, sede Concepción, Chile. *Rev Chil Nutr.* 2008;35(3):215–24.
23. 2Alzate-Ramirez J, Osorio-Quintero Y. Evaluacion de los habitos fisicos y saludables en estudiantes universitarios. Tesis Licenciatura Universidad del Valle Santiago Cali Colomb. 2013;1–86.
24. Vargas-Zárate M, Becerra-Bulla F, Prieto-Suárez E. Evaluación de la ingesta dietética en estudiantes universitarios. Bogotá, Colombia. *Rev Salud Publica.* 2010;12(1):116–25.
25. Guenther PM, Dodd KW, Reedy J, Krebs-Smith SM. A figure is presented most Americans Eat Much Less than Recommended Amounts of Fruits and Vegetables. *J Am Diet Assoc.* 2006;106(9): 1371–9.
26. Góme-Gutiérrez LF, Lucumí Cuesta DI, Girón Vargas SL, Espinosa-García G. Conglomeración de factores de riesgo de comportamiento asociados a enfermedades crónicas en adultos jóvenes de dos localidades de Bogotá, Colombia: Importancia de las diferencias de género. *Rev Esp Salud Publica.* 2004;78(4):493–504.
27. Malm C, Jakobsson J, Isaksson A. Physical Activity and Sports—Real Health Benefits: A Review with Insight into the Public Health of Sweden. *Sports.* 2019;7(5):127.
28. Ha EJ, Caine-Bish N. Effect of Nutrition Intervention Using a General Nutrition Course for Promoting Fruit and Vegetable Consumption among College Students. *J Nutr Educ Behav.* 2009;41(2):103–9.
29. Calfas KJ, Sallis JF, Nichols JF, Sarkin JA, Johnson MF, Caparosa S, et al. Project GRAD: Two-year outcomes of a randomized controlled physical activity intervention among young adults. *Am J Prev Med.* 2000;18(1):28–37.
30. Kahn EB, Ramsey LT, Brownson RC, Heath GW, Howze EH, Powell KE, et al. The Effectiveness of Interventions to Increase Physical Activity A Systematic Review. *Am J Prev Med.* 2010;22(02):73–107.